Availa® Cr 3%

A nutritional feed ingredient for animal feed. When used as a commercial feed ingredient, it must be declared as chromium methionine complex.



TYPICAL ANALYSIS

| MINERALS | Chromium | 30,000 ppm |
|-------------|---------------|------------|
| | | (3.0%) |
| | | |
| PROXIMATES | Moisture | 0.5 - 2.5% |
| | Crude Protein | 16.0% |
| | | |
| AMINO ACIDS | Methionine | 22.0% |

PHYSICAL CHARACTERISTICS

| Color | Purple |
|-----------------------------|------------------------------------|
| Texture | Granular |
| Particle Characteristics | Free flowing, slightly hygroscopic |
| Density | 480 - 540 kg/m ³ |
| Solubility | Active product is water soluble |
| Package Type | Multiwall bag |
| Package Weight | 20 kg |

SAFE HANDLING & STORAGE

When correctly used, there is no toxicity hazard in the use of Availa® Cr 3%. Refer to the SDS for more detail. Store in a clean and dry environment. Follow label directions and rotate inventory to ensure fresh product.

FEEDING INSTRUCTIONS

| DAIRY CATTLE | Non-Lactating Cows 0.55 PPM of mineral | |
|--------------|--|--|
| | Lactating Cows | 0.35 PPM of mineral |
| BEEF CATTLE | Weaned Calves | 0.4 PPM of mineral |
| | Cows and Bulls | 0.14 g/head/day |
| | Feedlot Receiving | 0.4 PPM of mineral |
| | Feedlot Finishing | 0.2 PPM of mineral |
| SHEEP | All | Feed at 0.015 g/50 kg body weight. Not recommended for use outside opremixes. |
| GOATS | All | Feed at 0.015 g/50 kg body weight. Not recommended for use outside opremixes. |
| HORSES | All | Feed at 0.03 g/100 kg body weight. Not recommended for use outside opremixes. |
| SWINE | Nursery | Up to 15 g/MT |
| | Grow/Finish | Up to 15 g/MT |
| | Sows and Boars | Up to 15 g/MT |
| POULTRY | All | 10 g/MT to reach 0.3 PPM in the complete feed |
| AQUACULTURE | Salmonids and Marine Fish | 135 g/MT |
| | Tilapia and Freshwater Fish | 135 g/MT |

Make sure adequate feed is always available to cows consuming Availa Cr 3%. Not recommended for use outside of premixes.