



## PRODUCT INFORMATION SHEET

# Perform-Max™ OPTIMIZER®

Water acidifier to be used through water proportioning systems for growing and finishing swine, broiler chickens and turkeys.



## PHYSICAL CHARACTERISTICS

<b>Color</b>	Blue Green
<b>Form</b>	Liquid
<b>Density</b>	.30 kg/L
<b>Solubility</b>	Soluble in Water
<b>Package Type</b>	4 X 1 Gallon Jugs
<b>Package Weight</b>	3.93 kg
<b>Volume</b>	128 fl oz (3.785 L)

## SAFE HANDLING & STORAGE

Store in a dark, cool, dry room (between 40 – 80°F [4 – 26°C]). Do not allow it to freeze. Do not leave the container open for long periods of time. Use only as according to label directions. Do not mix Zinpro® Specialty Products® with other nutritional supplements or chemicals without first testing compatibility. Manufacturer accepts no liability for improper use, handling, or storage. For oral use only. Keep out of reach of children. Not for human consumption.”

## DIRECTIONS FOR USE

### SWINE

#### All

3 days prior to harvest of finished market swine: Meter undiluted at a rate of 1 fl oz (29.57 ml) per gallon (per 3.78 liters) of finished drinking water (1:128) the last 3 days prior to harvest.

During periods of decreased feed and water consumption for growing swine: Provide Perform-Max Optimizer® up to 5 days per week. Mix 1 quart (.95 liter) in 2 gallons (7.57 liters) of stock solution and meter at a rate of 1 fl oz (29.57 ml) per gallon (per 3.78 liters) of finished drinking water (1:128).

### POULTRY

#### Broilers

3 days prior to harvest of finished market broilers: Meter undiluted at a rate of 1 fl oz (29.57 ml) per gallon (per 3.78 liters) of finished drinking water (1:128) the last 3 days prior to harvest.

During periods of decreased feed and water consumption for growing broiler chickens: Provide Perform-Max Optimizer® up to 5 days per week. Mix 1 quart (.95 liter) in 2 gallons (7.57 liters) of stock solution and meter at a rate of 1 fl oz (29.57 ml) per gallon (per 3.78 liters) of finished drinking water (1:128).

#### Turkeys

3 days prior to harvest of finished market turkeys: Meter undiluted at a rate of 1 fl oz (29.57 ml) per gallon (per 3.78 liters) of finished drinking water (1:128) the last 3 days prior to harvest.

During periods of decreased feed and water consumption for growing turkeys: Provide Perform-Max Optimizer® up to 5 days per week. Mix 1 quart (.95 liter) in 2 gallons (7.57 liters) of stock solution and meter at a rate of 1 fl oz (29.57 ml) per gallon (per 3.78 liters) of finished drinking water (1:128).

May be run in conjunction with Zinpro Performance Minerals® programs for optimal feed and water nutrient absorption and utilization.